Genome-wide association study of intraocular pressure uncovers new pathways to glaucoma

MacGregor S (1), Ong JS (2), An J (2), Han X (2), Zhou T (3), Siggs OM (3), Law MH (2), Souzeau E (3), Sharma S (3), Lynn DJ (4,5,6), Beesley J (2), Sheldrick B (3), Mills RA (3), Landers J (3), Ruddle JB (7), Graham SL (8), Healey PR (9,10), White AJR (9), Casson RJ (11), Best S (12), Grigg JR (10), Goldberg I (10), Powell JE (13,14), Whiteman DC (2), Radford-Smith GL (2,15), Martin NG (2), Montgomery GW (16), Burdon KP (3,17), Mackey DA (17,18), Gharahkhani P (2), Craig JE (3), Hewitt AW (7,17)

- 1 QIMR Berghofer Medical Research Institute, Brisbane, Queensland, Australia. stuart.macgregor@qimrberghofer.edu.au.
- 2 QIMR Berghofer Medical Research Institute, Brisbane, Queensland, Australia.
- 3 Department of Ophthalmology, Flinders University, Flinders Medical Centre, Bedford Park, South Australia, Australia.
- 4 South Australian Health and Medical Research Institute, Adelaide, South Australia, Australia.
- 5 EMBL Australia Group, Infection & Immunity Theme, South Australian Health and Medical Research Institute, Adelaide, South Australia, Australia.
- 6 College of Medicine and Public Health, Flinders University, Bedford Park, South Australia, Australia.
- 7 Centre for Eye Research Australia, University of Melbourne, Melbourne, Victoria, Australia.
- 8 Faculty of Medicine and Health Sciences, Macquarie University, Sydney, New South Wales, Australia.
- 9 Centre for Vision Research, Westmead Institute for Medical Research, University of Sydney, Sydney, New South Wales, Australia.
- 10 Discipline of Ophthalmology, Faculty of Medicine & Health Sciences, University of Sydney, Sydney Eye Hospital, Sydney, New South Wales, Australia.
- 11 South Australian Institute of Ophthalmology, University of Adelaide, Adelaide, South Australia, Australia.
- 12 Eye Department, Greenlane Clinical Centre, Auckland District Health Board, Auckland, New Zealand.
- 13 Garvan-Weizmann Centre for Cellular Genomics, Garvan Institute of Medical Research, Sydney, New South Wales, Australia.
- 14 St Vincent's Clinical School, University of New South Wales, Sydney, New South Wales, Australia.
- 15 University of Queensland School of Medicine, Brisbane, Queensland, Australia.
- 16 Institute for Molecular Bioscience, University of Queensland, Brisbane, Queensland, Australia.
- 17 Menzies Institute for Medical Research, University of Tasmania, Hobart, Tasmania, Australia.
- 18 Centre for Ophthalmology and Visual Science, University of Western Australia, Crawley, Western Australia, Australia.

Intraocular pressure (IOP) is currently the sole modifiable risk factor for primary open-angle glaucoma (POAG), one of the leading causes of blindness worldwide1. Both IOP and POAG are highly heritable2. We report a combined analysis of participants from the UK Biobank (n?=?103,914) and previously published data from the UK Biobank (n?=?103,914).

Nat Genet. 2018 Aug;50(8):1067-1071. doi: 10.1038/s41588-018-0176-y. Epub 2018 Jul 27.

http://www.ncbi.nlm.nih.gov/pubmed/30054594